

Immunogenicity and Immunomodulatory Properties of Umbilical Cord Lining Mesenchymal Stem Cells

Tobias Deuse,*†‡ Mandy Stubbendorff,† Karis Tang-Quan,†‡ Neil Phillips,§
Mark A. Kay,§ Thomas Eiermann,¶ Thang T. Phan,##* Hans-Dieter Volk,††
Hermann Reichenspurner,*† Robert C. Robbins,‡ and Sonja Schrepfer*†‡

*Cardiovascular Surgery, University Heart Center Hamburg, Hamburg, Germany

†TSI-lab, Cardiovascular Research Center, University Hospital Hamburg-Eppendorf, Hamburg, Germany

‡Cardiothoracic Surgery, Stanford University School of Medicine, Stanford, CA, USA

§Pediatrics, Human Gene Therapy, Stanford University School of Medicine, Stanford, CA, USA

¶Transfusion Medicine, HLA-Lab, University Hospital Hamburg-Eppendorf, Hamburg, Germany

#Surgery, Yong Loo Lin School of Medicine, Singapore

*CellResearchCorp, Clinical Research Centre, Singapore

††BCRT, Charite, Campus Virchow-Klinikum, Berlin, Germany

We here present an immunologic head-to-head comparison between human umbilical cord lining mesenchymal stem cells (cMSCs) and adult bone marrow MSCs (bmMSCs) from patients >65 years of age. cMSCs had significantly lower HLA class I expression, higher production of tolerogenic TGF- β and IL-10, and showed significantly faster proliferation. In vitro activation of allogeneic lymphocytes and xenogeneic in vivo immune activation was significantly stronger with bmMSCs, whereas immune recognition of cMSCs was significantly weaker. Thus, bmMSCs were more quickly rejected in immunocompetent mice. IFN- γ at 25 ng/ml increased both immunogenicity by upregulation of HLA class I/ HLA-DR expression and tolerogenicity by increasing intracellular HLA-G and surface HLA-E expression, augmenting TGF- β and IL-10 release, and inducing indoleamine 2,3-dioxygenase (IDO) expression. Higher concentrations of IFN- γ (>50 ng/ml) further enhanced the immunosuppressive phenotype of cMSCs, more strongly downregulating HLA-DR expression and further increasing IDO production (at 500 ng/ml). The net functional immunosuppressive efficacy of MSCs was tested in mixed lymphocyte cultures. Although both cMSCs and bmMSCs significantly reduced in vitro immune activation, cMSCs were significantly more effective than bmMSCs. The veto function of both MSC lines was enhanced in escalating IFN- γ environments. In conclusion, cMSCs show a more beneficial immunogenic profile and stronger overall immunosuppressive potential than aged bmMSCs.

Key words: Mesenchymal stem cells (MSCs); Immunogenicity; Immunomodulation; Stem cell transplantation

INTRODUCTION

Mesenchymal stem cells (MSCs) have been identified a promising cell population not only for tissue regeneration (15,30,40), but also for immunomodulatory therapy (31,42). An emerging body of data indicates that MSCs may play specific roles as immunomodulators in maintenance of peripheral tolerance, transplantation tolerance, autoimmunity, and tumor immunobiology (25). MSCs were shown to interact with various immune cell types including T cells (9,18,41), B cells (1,7,11), dendritic cells (DCs) (14,48), and natural killer (NK) cells (17,37). Mul-

tiple mechanisms, requiring dynamic cross-talk between MSCs and immune cells and involving soluble immunomodulatory factors, cause a shift of the immune response towards tolerance or anergy (11,28,46). Release of transforming growth factor- β (TGF- β), hepatocyte growth factor (HGF), and induction of the tryptophan catabolizing enzyme indoleamine 2,3-dioxygenase (IDO) has been suggested to mediate the immunosuppressive effects of MSCs (9,22).

The immunosuppressive properties of MSCs have been examined in a variety of animal models of autoimmunity, solid organ, or stem cell transplantation. MSCs

Received April 8, 2010; final acceptance September 22, 2010. Online prepub date: November 5, 2010.

Address correspondence to Sonja Schrepfer, M.D., Ph.D., Heisenbergprofessor at the University Heart Center Hamburg, Transplant and Stem Cell Immunobiology Lab (TSI), Martinistr. 52, Campus Research, N27, Room 068, 20146 Hamburg, Germany. Tel: +49-40-7410-5 9982; Fax: +49-40-7410-5 9663; E-mail: schrepfer@stanford.edu

have been reported to ameliorate experimental autoimmune encephalomyelitis (46,47) in mice. Infusion of host-derived MSCs has been shown to decrease the rejection of allogeneic bone marrow (BM) stem cells in a murine transplant model (26). Even more impressive, MSC administration sufficiently reversed ongoing graft versus host disease (GVHD) (45) in mice, an effect that could already be verified in human disease (19,33). Modest success in prolonging the survival of allogeneic skin transplants by MSC infusion has already been demonstrated in baboons (3). However, controversies exist on the efficiency of MSC infusion to prolong solid organ graft survival (13,50). The involved mechanisms of MSC-based immunomodulation may favor the suppression of a more generalized state of immune activation in BM transplant, GVHD, or autoimmune disease compared to the more localized immune response in organ transplantation (2). Still, recommendations have been formulated on how to move MSC-based tolerance strategies forward toward clinical application in solid organ transplantation (8).

Because MSCs can be isolated from various tissues including BM, adipose tissue (51), umbilical cord (34), placenta, liver, and spleen (4), and these cell populations may vary in their functional capacities, the optimal source of MSCs for cell-based immunomodulatory therapies remains to be determined. Patients in need of stem cell-based therapies for cardiovascular diseases are usually aged and approaches using autologous adult stem cells have to deal with aged cells. Allogeneic stem cell strategies can utilize fetal stem cells, but may encounter problems with immune activation and cell rejection. We here present an immunologic head-to-head comparison between human umbilical cord lining MSCs (cMSCs) and aged adult BM MSCs (bmMSCs) to help clarify which strategy offers better immunomodulatory properties.

MATERIALS AND METHODS

Animals

Six- to 8-week-old male Balb/c and Scid-beige (CB17.Cg-Prkdc^{scid} Lyst^{tg}/CrI) mice were purchased from Charles River Laboratories (Sulzfeld, Germany) and received humane care in compliance with the Guide for the Principles of Laboratory Animals.

MSC Culture and Cell Transplantation

Female neonatal cMSCs (kindly provided by Cell Research Corporation Pte Ltd, Singapore) were isolated from the outer amniotic lining of the umbilical cord and were cultured in PTT4 media (CellResearch). bmMSCs were isolated from the bone marrow of three different healthy female donors more than 65 years of age (Promocell, Heidelberg). Therefore, the bone marrow was

flushed from femoral heads and isolated using plastic adherence. Cells were grown in their specific growth media [CMRL-1066 + antibiotics + L-glutamine (Cell ResearchCorp) for cMSCs and mesenchymal stem cell growth media (Promocell) for bmMSCs]. Cells were split at a confluency of 80–85% by mechanically lifting the cells from the tissue culture plastic surface using a cell lifter (Costar, Corning). Before transplantation, MSCs were dissociated by collagenase type I solution at a concentration of 0.025% for 5 min at 37°C, and resuspended in sterile PBS at 1×10^6 cells per 50 μ l. MSC viability was approximately 95% as determined by trypan blue staining. MSC transplantation was performed by direct injection of 1×10^6 MSCs into the right gastrocnemius muscle of recipient mice using a 27-gauge syringe. In vitro experiments with MSCs were also performed at a confluency of 80–85%.

Multipotent Capacity

Adipogenic Differentiation. Cells were plated onto 24-well plates at 6×10^4 cells per well (3.15×10^4 cells/cm²) and differentiation was induced at 80–90% confluency by using mesenchymal stem cell adipogenic differentiation medium (Promocell) for 14 days at 37°C and 5% CO₂. Medium was changed every third day. After 2 weeks, cells were washed with PBS (Gibco) and fixed using 10% formalin for 30 min at room temperature. The Oil Red O stock solution was freshly prepared by dissolving 300 mg Oil Red O in 100 ml 99% isopropanol. Fixed cells were washed with distilled water (Gibco), incubated with isopropanol for 5 min followed by incubation with filtered Oil Red O working solution for 5 min. Hematoxylin stain was added for 30 s, followed by 1% acetic acid. Lipids appear red and nuclei appear blue on a phase contrast microscope.

Chondrogenic Differentiation. Cells (5×10^5) were plated into a 96-well U-bottom suspension culture plate using ready-to-use mesenchymal stem cell chondrogenic differentiation medium (Promocell) in an incubator with an atmosphere of 5% CO₂ at 37°C. Spheroids spontaneously formed within 24–48 h. Cells were incubated for 21 days; medium was changed every third day without aspirating the spheroids.

Spheroids were placed into 10% formalin for 1 h, dehydrated, and paraffin embedded. Paraffin sections (5 μ m) were prepared, and slides were deparaffinized, hydrated, and stained with toluidine blue (1 g toluidine blue in 100 ml 70% ethanol) for 3 min. Slides were dehydrated through 95% and 100% ethanol, and cleared in xylene. Chondrogenic spheroids appear blue on a phase contrast microscope.

Osteogenic Differentiation. Cells (6×10^4) were plated onto a 24-well tissue culture plate (3.15×10^4 cells/cm²) and differentiation was induced after cells

reached >100% confluency (after 24–72 h) in an incubator at 37°C and 5% CO₂. Cells were incubated with mesenchymal stem cell osteogenic differentiation medium (Promocell) for 21 days; medium was changed every third day. Cells were fixed by removing the media, washing with PBS, and incubating with 10% formalin for 10 min. Alizarin Red S solution (2 g Alizarin red S in 100 ml dest. water) was incubated for 2 min. The osteogenic lineage was identified by dark red staining on a phase contrast microscope.

MSC Proliferation Assay

MSCs were seeded in 24-well plates (3,000 cells/well) and grown in their specific growth media. Daily cell counts were determined using the CellTiter 96 Cell Aqueous One Solution Proliferation Assay (Promega, Madison, WI). Triplicates were performed and the absorbance at 490 nm was measured with the Magellan ELISA Reader and Software (Tecan Systems Inc, San Jose, CA).

TGF- β and IL-10 Elispot Assays

The cytokine release for TGF- β and IL-10 was analyzed using human Elispot assays (R&D Systems). cIMSCs or bmMSCs (5×10^5) were plated in their cell-specific growth media onto TGF- β and IL-10-coated Elispot plates for 48 h in an incubator at 37°C and 5% CO₂.

Spot developing was performed according to the manufacturer's protocol (BD Biosciences). Spots were automatically enumerated using an Elispot plate reader (CTL, OH) for scanning and analyzing.

IDO Western Blots

Cell extract was prepared in RIPA buffer (Sigma). Amount of protein was determined and electrophoresis was performed by loading an 8 μ g protein sample into each well. Proteins were then blotted onto PVDF membrane in transfer buffer (3.75 g glycine, 7.25 g Tris base, 2.5 ml methanol was dissolved in total 2.5 L). The membranes were incubated in 5% BSA/TPBS solution prepared in PBST buffer (1 $\times$ PBS, pH 7.0, 1 ml Tween-20/L) at room temperature for 2 h and then primary antibodies were added overnight at 4°C in 1:5000 dilution for IDO. Once the primary antibody labeling was performed, the blots were washed with PBST and the secondary antibody labeling (goat anti-mouse conjugated to horseradish peroxidase) was performed at room temperature for 1 h at 1:5000 dilution in 5% BSA in PBST. After 1 h of washing at room temperature with PBST, the ECL Western Blot System (Amersham, Piscataway, NJ) was used to visualize proteins using bioluminescence imaging (Caliper, IVIS 200), and photons per second were measured. Membrane-bound antibodies were

stripped using restore Western blot stripping buffer (Pierce) for 15 min at room temperature. Membranes were blocked for 2 h using in 5% BSA/TPBS buffer, and primary antibody for GAPDH was added overnight at 4°C.

Bioluminescence Imaging (BLI)

BLI is a noninvasive, molecular imaging modality that allows for quantitative and repetitive imaging of targeted cells, transfected with plasmid vectors encoding for firefly luciferase (Fluc), in living organisms (44). In the presence of ATP and O₂, Fluc converts exogenously administered D-luciferin to the optically active metabolite oxyluciferin. Light emission can be monitored with an ultrasensitive CCD camera. In the setting of cell transplantation, BLI signals correlate with the number of living Fluc⁺ cells. For this study, MSCs were transduced with Adv-Fluc (CMV) at a multiplicity of infection of 10 to stably express Fluc. Briefly, after MSCs were seeded, viral aliquots were added to 0.75 ml of their cell-specific growth media containing 8 μ g/ml polybrene at a multiplicity of infection of 10, and the plate was then placed into a cell incubator at 37°C and 5% CO₂ for 24 h. Medium was changed after 24 h. Transduction efficacy was confirmed using Adv-GFP.

BLI was performed using the Xenogen in vivo imaging system (Caliper Life Sciences, Hopkinton, MA) as reported previously (5). Mice were anesthetized with 2% isoflurane and D-luciferin (Caliper Life Sciences) was administered IP at a dose of 375 mg/kg of body weight. At the time of imaging, animals were placed in a light-tight chamber, and photons emitted from FLuc-expressing MSCs were counted. The recipient mice were scanned repetitively on days 0, 1, and then every other day until the signal dropped into the background. BLI signals were quantified in units of photons per second (total flux) and are presented on a logarithmic scale.

Cell Surface Molecule Expression

MSCs grown in their growth media with or without additional recombinant human IFN- γ at 25 ng/ml (low-dose; PeproTech) were harvested to assess their expression of various surface markers. Cells were dissociated by collagenase type I solution at a concentration of 0.025% for 5 min at 37°C, and 5×10^5 cells were stained for 45 min at 4°C in 100 μ l of 0.1% BSA in PBS containing an appropriate dilution of a desired PE-conjugated antibody. Primary PE-conjugated antibodies (all BD Bioscience unless stated otherwise) used were anti-human-CD31 (clone WM59), CD44 (clone G44-26), CD34 (clone 563), CD45 (clone H130), CD90 (clone 5E10), CD105 (clone 166707), CD117 (clone 104D2), SSEA-1 (clone MC-480), SSEA-4 (clone MC813-70), TRA-1-60 (clone TRA-1-60), HLA class I (clone

DX17), β_2 -microglobulin (clone T $\ddot{U}$ 99), HLA-DR (clone WR18; abcam), HLA-G (clone MEM-G/9; Santa Cruz), HLA-E (clone MEM-E/06; Santa Cruz), CD40 (clone 5C3), CD80 (clone L307.4), and CD86 (clone 2331). IgG3 (clone 133316) isotype control was purchased from R&D Systems (Minneapolis, MN), IgG2a isotype control from Abcam (clone MOPC-173; Cambridge, MA), IgG1 (clone MOPC-21), and IgM (clone G155-228) from BD Biosciences. For the detection of intracellular HLA-G, cells were made permeable using the fix and perm cell permeabilization kit (BD Bioscience).

Flow cytometry was performed on a FACSCalibur system with Cell Quest Pro software (Becton Dickinson, Heidelberg, Germany); 10,000 events were collected. Analysis of data was performed using Flowjo (Tree Star, Ashland, OR). In the histograms, the isotype controls are represented by the gray solid areas and the antigens of interest by solid lines. Mean fluorescence intensities (MFI) were calculated. Molecule expression is defined as the ratio (MFI of antigen)/(MFI of negative isotype control), with 1 equaling no expression.

Elispot Assays

The cellular in vivo immune responses in Balb/c mice were studied after cell transplantation of either cMSCs or bmMSCs. The spleen of recipient animals was harvested 5 days after MSC transplantation to isolate recipient splenocytes. Elispot assays using 5×10^5 mitomycin-inhibited MSCs (Sigma Aldrich, St. Louis, MO) as stimulator cells and 5×10^6 recipient splenocytes as responder cells were performed according to the manufacturer's protocol (BD Biosciences) using IFN- γ and IL-4-coated plates.

The immunogenicity of MSCs to elicit an in vitro immune response was investigated by incubating 1×10^5 cMSCs or bmMSCs with 1×10^6 human peripheral blood mononuclear cells (PBMCs) for 7 days on a coated Elispot plate. Human PBMCs from healthy donors were previously isolated by Ficoll gradient centrifugation.

The immunosuppressive properties of MSCs in vitro were assessed by their potential to suppress a one-directional allogeneic mixed lymphocyte response in an allogeneic setting. Mitomycin-inhibited stimulator lymphocytes (1×10^5 , freshly isolated from healthy donors by Ficoll) and 1×10^6 allogeneic effector lymphocytes were incubated with 1×10^5 cMSCs or bmMSCs in an IFN- γ , IL-4, or IL-2-coated Elispot plate for 7 days. Plates were developed according to the manufacturer's protocol (BD Bioscience) and spots were automatically enumerated using an Elispot plate reader (CTL, OH) for scanning and analyzing.

For IL-2 Elislots, cMSCs and bmMSCs were used

either untreated or pretreated with 25 or 500 ng/ml IFN- γ for 48 h.

The High-IFN- γ Veto Function of MSCs

MSCs grown in their growth media with or without additional recombinant human IFN- γ at 25 ng/ml (low dose) or 500 ng/ml (high dose). Cells were harvested to assess their surface expression of HLA class I, HLA-DR, HLA-G, and HLA-E by flow cytometry as described above. Also, their IDO production was quantified in Western blot assays.

Statistical Analysis

Data are presented as mean $\pm$ SD. Comparisons between two groups were done by independent sample *t*-tests. Comparisons between groups were done by analysis of variance between groups (ANOVA) with LSD Post Hoc tests. Differences were considered significant for $p < 0.05$. Statistical analysis was performed using SPSS statistical software for Windows (SPSS, Chicago, IL).

RESULTS

Phenotypic and Functional Characteristics of MSCs

Both cMSCs and bmMSCs were positive for the mesenchymal stem cell markers CD90 and CD105 and negative for hematopoietic stem cell markers such as CD34, CD45, and CD117 (Fig. 1A). They were positive for the cell-cell interaction protein CD44 and negative for CD31. Both MSC lines were mildly positive for SSEA-4, an embryonic stem cell marker, and remained negative for SSEA-1, a marker for maturity, and the keratan sulfate antigen TRA-1-60, a human embryonic carcinoma marker.

In culture, both cMSCs and bmMSCs had fibroblast-like spindle-shaped morphology (Fig. 1B), although cMSCs were mildly bigger. After trypsination, the rounded cMSCs had diameters of $15.8 \pm 3.1 \mu\text{m}$ and the bmMSCs of $11.8 \pm 4.1 \mu\text{m}$ ($p < 0.001$). Both MSC lines were easily differentiable towards the chondrogenic, adipogenic, and osteogenic lineages. No differences between both MSC lines were noted regarding the time required for cell differentiation.

When grown in their cell specific media, cell proliferation of cMSCs was much faster than of bmMSCs (Fig. 1C). Differences between groups were statistically significant ($p < 0.05$) for day 3 and later time points. Thus, cell cultures with cMSCs required splitting approximately every 3 days, whereas bmMSCs were split on average every 10 days.

Antigenic, Costimulatory, and Tolerogenic Properties of MSCs

MSCs were screened for their surface expression of HLA antigens, costimulatory factors, and immune toler-




Figure 1. Phenotypic and functional characteristics of MSCs. cIMSCs and bmMSCs were positive for CD90, CD105, and CD44, mildly positive for SSEA-4, and negative for CD31, CD34, CD45, CD117, SSEA-1, and TRA-1-60 (A; isotype control: gray solid area, antigens of interest: solid line; molecule expression is provided in the right upper corners). Both MSC lines grew in fibroblast-like spindle shape during cell culture conditions (original magnification 200×) and were differentiable into chondrocytes (original magnification 100×), adipocytes (original magnification 200×), and osteocytes (original magnification 100×, Leica microscope with QWin acquisition software, B). In vitro, cIMSCs proliferation over a 5-day period was much faster than proliferation of bmMSCs (C; * $p < 0.05$, † $p < 0.01$).

ance molecules at rest and in an inflammatory milieu with IFN- γ at 25 ng/ml. cIMSCs had far lower expressions of HLA class I molecules ($p < 0.001$) and β_2 -microglobulin ($p = 0.001$) than bmMSCs. Both MSCs completely lacked HLA-DR expression and were negative for the costimulatory factors CD40, CD80, and CD86 (Fig. 2A). After stimulation with IFN- γ , the levels of HLA class I ($p = 0.014$ and $p = 0.015$) and HLA-DR ($p < 0.001$ each) significantly increased for cIMSCs and bmMSCs, respectively. All costimulatory factors remained negative after IFN- γ challenge. Interestingly,

surface expression of HLA-E ($p = 0.012$ for cIMSCs and $p = 0.033$ for bmMSCs), but not HLA-G (not shown) was significantly upregulated with IFN- γ in both MSC lines. Intracellular HLA-G, however, was constitutively expressed in both MSCs, although at higher levels in cIMSCs ($p = 0.034$). HLA-G was upregulated by IFN- γ ($p = 0.005$ for cIMSCs and $p = 0.043$ for bmMSCs).

The release of the tolerogenic cytokines TGF- β and IL-10 was quantified in Elispot assays. At rest, both MSC lines produced TGF- β , but only cIMSCs released detectable amounts of IL-10 (Fig. 2B). After IFN- γ stim-




Figure 2. Antigenic, costimulatory, and tolerogenic properties of MSCs. Surface molecule expression of cIMSCs and bmMSCs was assessed by flow cytometry in the presence and absence of IFN- γ at 25 ng/ml (A; * p < 0.05, † p < 0.01). TGF- β and IL-10 production by MSCs was quantified in ELISPOT assays (B; * p < 0.05, † p < 0.01). With and without IFN- γ , the cytokine release was significantly higher in cIMSC cultures. The enzyme IDO was hardly detectable in resting MSCs using Western blots, but IDO expression was strongly upregulated by IFN- γ in both cIMSCs and bmMSCs (C; * p < 0.05, † p < 0.01).

ulation, the release of both TGF- β (p = 0.018 and p = 0.003) and IL-10 (p < 0.001 and p < 0.001) was significantly increased by cIMSCs and bmMSCs, respectively. Absolute values for TGF- β and IL-10 release were significantly higher in cIMSC cultures (p = 0.03 and p < 0.001, respectively) than with bmMSCs. IDO expression was hardly detectable in resting MSCs, but was strongly upregulated by IFN- γ in cIMSCs (p < 0.001) and bmMSCs (p = 0.041) (Fig. 2C).

Immunogenicity of MSCs

Because cIMSCs and bmMSCs were shown to have differences in their expression of immunogenic molecules as well as in their release of tolerogenic factors, it was tested if they were immunogenic enough to elicit an immune response in vivo and in vitro. In preparation for BLI, cIMSCs and bmMSCs tolerated the FLuc transduction well and both showed a high transduction efficacy

of >95% as confirmed by Adv-GFP/DAPI double staining (not shown).

Elispot assays, performed 5 days after xenogeneic MSC transplantation, showed that both MSC lines triggered some cellular immune activation. However, bmMSCs were much more immunogenic and the provoked Th1 and Th2 responses were much stronger. Spot frequencies for IFN- γ and IL-4 were 10.8-fold (95% CI: 6.6–17.8) and 3.4-fold (95% CI: 2.2–5.2) higher with bmMSCs, respectively ($p < 0.001$ each vs. cMSCs) (Fig. 3A). Concordantly, bmMSCs were rejected more quickly than cMSCs in immunocompetent Balb/c mice. BLI signals dropped into the background after 7.2 ± 0.9 days for bmMSCs and after 10.9 ± 1.2 days for cMSCs ($p < 0.001$) (Fig. 3B). In immunodeficient SCID-beige mice, cell survival was prolonged and similar (14.7 ± 2.5 and 17.0 ± 2.8 days, respectively, $p = 0.205$).

Allogeneic immune activation was assessed in Elispot assays after incubation of MSCs with human PBMCs. Both MSC lines showed five out of six possible HLA-A, HLA-B, and HLA-DR mismatches with the used effector lymphocytes, but all cells shared the HLA-A24 epitope (Table 1). We found only very weak in vitro T-cell activation by cMSCs (PBMC + cMSC) with non-significant increases in IFN- γ spots (ANOVA $p = 0.002$, LSD post hoc $p = 0.419$) and IL-2 spots (ANOVA $p = 0.042$, LSD post hoc $p = 0.060$) versus PBMC control. Allogeneic in vitro lymphocyte activation, however, was stronger for bmMSCs (PBMC + bmMSC) with significantly increased IFN- γ (LSD post hoc $p = 0.001$) and IL-2 spots (LSD post hoc $p = 0.014$) (Fig. 3C) versus PBMC.

These data demonstrate that although both cMSCs and bmMSCs are recognized by allogeneic and xenogeneic lymphocytes, cMSCs are less immunogenic, cause less immune activation, and get rejected more slowly.

Immunosuppressive Properties of MSCs

HLA molecule expression of MSCs was further assessed in escalating IFN- γ environments. Although both cMSCs and bmMSCs upregulated HLA-DR expression upon low-dose IFN- γ stimulation at 25–50 ng/ml, HLA-DR was downregulated again with higher doses (100–500 ng/ml) (Fig. 4A). Throughout the IFN- γ range of 25–500 ng/ml, HLA-DR expression and HLA-I expression were significantly lower in cMSCs compared to bmMSCs ($p \leq 0.020$). Although HLA-I expressions also showed mild reductions with increasing IFN- γ concentrations, the downregulation remained statistically insignificant for both cMSCs and bmMSCs. Intracellular HLA-G showed a mild increase with rising IFN- γ that reached significance for the 500 ng/ml concentration compared to unstimulated conditions in both cell lines.

HLA-E surface expression was already significantly upregulated by 25 ng/ml IFN- γ and remained at similar levels in higher IFN- γ milieus.

With increasing IFN- γ concentrations from 25 to 500 ng/ml, IDO expression further significantly increased from 1.2 ± 0.1 to 2.3 ± 0.1 ($p < 0.001$) in cMSCs (Fig. 4B). No significant differences were observed between 25 and 100 ng/ml IFN- γ in cMSCs and throughout the IFN- γ range in bmMSCs.

To test their net immunosuppressive effects, MSCs were incubated with allogeneic mixed PBMC cultures at different IFN- γ concentrations. Stimulator and effector lymphocytes had three out of four possible HLA-A/B mismatches and one out of two possible HLA-DR mismatches (Table 1). Effector lymphocyte activation was assessed by IL-2 Elispot assays (Fig. 4C). Both cMSCs ($p < 0.001$) and bmMSCs ($p < 0.001$) significantly reduced the IL-2 spot frequencies of mixed PBMC cultures in the absence of IFN- γ , although cMSCs were significantly more potent than bmMSCs ($p < 0.001$). Efficacy of bmMSCs was significantly improved in a low-dose IFN- γ environment (25 ng/ml; $p < 0.001$) and IL-2 spots were significantly lower with 500 versus 25 ng/ml IFN- γ in both MSCs ($p < 0.001$ each).

DISCUSSION

There are fundamental biological and practical differences between cMSCs and bmMSCs for cell-based immunomodulation. Whereas bmMSCs can be individually tailored for autologous use in chronic diseases, the retrieval of sufficient amounts of multipotent cells from BM of aging patients becomes increasingly problematic. Individualized cell production and storage would be very costly. On the other side, cMSCs are abundantly available, and can easily be isolated and expanded for a commercial off-the-shelf product for allogeneic use. The effect of MSCs seems not to be donor restricted, because the survival of third-party grafts could also be prolonged and third-party MSCs have also been shown to alleviate the rejection of skin grafts in baboons (3). MSCs were shown to modulate mixed lymphocyte reactions independent of MHC matching status (20) and third-party MSCs were shown to facilitate engraftment after umbilical cord blood transplantation (16). Third-party MSC infusion in hematopoietic stem cell transplant patients did not induce immunological memory to the infused MSCs (39). These characteristics support the possibility of exploiting universal “one cell fits all” MSC lines for therapeutic applications.

Both cMSCs and bmMSCs used in our study met phenotypical and functional criteria required for the definition of MSCs (12,25). They express CD105, CD90, and CD44, and are devoid of hematopoietic and endo-




Figure 3. Immunogenicity of MSCs. Five days after MSC transplantation into Balb/c mice, Elispot assays for IFN- γ and IL-4 demonstrated that bmMSCs were much more immunogenic than cIMSCs (A; $\dagger p < 0.01$). Survival of FLuc⁺ MSCs in Balb/c mice and in SCID-beige mice was longitudinally followed by BLI (B). bmMSCs were rejected more quickly than cIMSCs in immunocompetent Balb/c mice, whereas MSC survival was similar and significantly longer in immunodeficient SCID-beige mice. Allogeneic immune activation in vitro after incubation of MSCs with human PBMCs showed very weak T-cell activation by cIMSCs but a significant lymphocyte activation by bmMSCs (C; $*p < 0.05$, $\dagger p < 0.01$).

thelial markers, such as CD31, CD34, CD45, and CD117. Because of the lack of specific MSC markers, additional antigens like the glycolipids SSEA-1 and SSEA-4 as well as the keratin sulfate TRA-1-60 are usually screened for. In line with previous reports (10,32), both of our MSC lines were positive for the embryonic stem cell-associated marker SSEA-4, although contrary findings were reported (36).

Additionally, cMSCs and bmMSCs allowed adipogenic, osteogenic, and chondrogenic differentiation *in vitro*. Neither cMSCs nor bmMSCs were positive for Oct4, Sox-2, or TRA-1-60 (data not shown), essential factors for maintenance of pluripotency and reflecting the primitiveness of embryonic stem cells (21,49). Even after serum deprivation, we and others (49) could not induce Oct4 expression, as has been reported earlier (27). In accordance with previous reports on osteogenic capacity of stem cells (49), we could show the far superior proliferation capacity of cMSCs over adult bmMSCs.

Our data show that MSCs may be considered hypo-immunogenic due to their relatively low MHC I expression, but they are not intrinsically immunoprivileged and undergo immune rejection in an immunocompetent host. The reduction in MSC survival times between immunodeficient SCID-beige mice and Balb/c mice must be attributed to host immune responses. We show that cMSCs express significantly lower levels of HLA-I at rest and after stimulation with IFN- γ . Recent evidence indicates that MSCs can function as antigen presenting cells (APC) themselves, with phagocytic function and the ability to induce T-cell responses (6,38). We demonstrate that, under resting cell culture conditions, both MSCs were negative for HLA-DR. In the presence of IFN- γ , however, HLA-DR was upregulated, especially in bmMSCs, in which significantly higher expression values were achieved compared to cMSCs. Because cell culture conditions can impact HLA-DR expression (35), all experiments were performed at similar cell densities. Both MSC lines were negative for costimulatory molecules. The lower HLA class I and II expression profile on cMSCs impedes immune recognition, promotes immune ignorance, and delays immune rejection *in vivo*. We found significantly longer survival times of cMSCs in Balb/c mice compared to bmMSCs. Also, xenogeneic *in vivo* immune activation of Th1 and Th2 cells upon MSC transplantation was weaker for cMSCs. In accordance with our xenotransplant results, allogeneic *in vitro* lymphocyte activation was mild for cMSCs and thus much weaker than for bmMSCs, which triggered considerable IFN- γ and IL-4 production. Although it has been reported that cMSCs from Wharton's jelly completely fail to induce allogeneic *in vitro* PBMC activation (43), we have used more sensitive Elispot assays and have

found at least mild immune activation with all MSC lines tested in our lab so far. However, MSCs gain tolerogenic potential through IDO expression (22), HLA-G production (24), and IL-10 and TGF- β secretion (23). These properties firstly alleviate immune recognition and diminish MSCs rejection and secondly may be used to suppress ongoing immune processes in the close vicinity of the administered or homed MSCs. We show that cMSCs produce larger amounts of tolerogenic IL-10 and TGF- β and have higher amounts of intracellular HLA-G.

IFN- γ dose-dependently exerts opposing effects on MSCs, enhancing their immunogenicity at low doses and increasing the production of tolerogenic cytokines and the expression of IDO at higher doses (6). It was previously shown that low-dose IFN- γ is produced by untreated MSCs, which may serve as endogenous autocrine stimulation (6) to enable their APC function. During inflammation, GVHD, and immune rejection, high levels of inflammatory mediators like IFN- γ are hypothesized to create a microenvironment in which the veto properties of MSCs dominate and MSCs suppress the ongoing immune reaction (6,29). We show that repression of HLA-DR is much more effective than of HLA-I in both cMSCs and bmMSCs. Head-to-head comparison between MSC lines shows that upregulated HLA-DR expression by 25 ng/ml IFN- γ remains much lower in cMSCs, and that HLA-DR downregulation by IFN- γ at concentrations beyond 50 ng/ml is also more effective in cMSCs. IDO production increases with rising IFN- γ , and shows no decrease at higher concentrations.

The net immunosuppressive effect of MSCs was tested in mixed PBMC cultures. The main findings are that cMSCs are more immunosuppressive than bmMSCs in the absence of IFN- γ and that the immunosuppressive effect of MSCs can be enhanced by IFN- γ . In an inflammatory milieu, MSCs seem to get further activated,

Table 1. Both MSC Lines Show Five Out of Six Possible HLA-A, HLA-B, and HLA-DR Mismatches With the Used Effector Lymphocytes, But All Cells Share the HLA-A24 Epitope

	HLA-A	HLA-B	HLA-DR
Effector lymphocytes	0301 2402	0702 —	1501 —
Stimulator lymphocytes	0101 2402	1501 4405	1501 1601
cMSCs	1101 2407	3505 3802	1101 1202
bmMSCs	0201 2402	3901 5101	1101 1301




Figure 4. Immunosuppressive properties of MSCs. An increase of the IFN- γ concentration beyond 50 ng/ml significantly downregulated cIMSC and bmMSC HLA-DR expression, whereas absolute HLA-DR values remained significantly lower in cIMSCs at all IFN- γ concentrations (A). IDO expression was significantly increased from 25 to 500 ng/ml IFN- γ in cIMSCs and moderately, nonsignificantly in bmMSCs (B). The net immunosuppressive effects of MSCs at different IFN- γ concentrations was assessed by in vitro IL-2 Elispot assays in allogeneic mixed PBMC cultures (C; S: stimulator PBMC, E: effector PBMC). cIMSCs were significantly more potent in reducing IL-2 spot frequencies. The efficacy of bmMSCs significantly improved with rising IFN- γ concentrations.

which enhances their immunosuppressive potential and makes them promising tools for future clinical use.

SUMMARY

We show that MSCs are not inherently immune evasive and immunoprivileged and that cMSCs and aged bmMSCs show relevant differences in their immunogenic profile and their capacity to respond to inflammatory stimuli and to suppress allogeneic immune activation. Altogether, cMSCs are superior cells for immunomodulation, because of their higher proliferative capacity, their lower immunogenicity, and their stronger production of soluble tolerogenic factors, contributing to their stronger overall immunosuppressive potential.

ACKNOWLEDGMENTS: We thank Christiane Pahrman (University Heart Center Hamburg, TSI-Lab) for her excellent assistance in performing the study, and Edwin Chow (Cell Research Corporation Pte Ltd) for providing cMSCs and media. Sonja Schrepfer received DFG Grants (SCHR992/3-1 and SCHR992/4-1). Tobias Deuse received the ISHLT Grant 2009. Karis Tang-Quan received the ESOT Short Stay Grant 2009. Thang T. Phan is affiliated with CellResearchCorp.; the other authors have no conflicts to disclose.

REFERENCES

- Augello, A.; Tasso, R.; Negrini, S. M.; Amateis, A.; Indiveri, F.; Cancedda, R.; Pennesi, G. Bone marrow mesenchymal progenitor cells inhibit lymphocyte proliferation by activation of the programmed death 1 pathway. *Eur. J. Immunol.* 35(5):1482–1490; 2005.
- Bartholomew, A.; Polchert, D.; Szilagyi, E.; Douglas, G. W.; Kenyon, N. Mesenchymal stem cells in the induction of transplantation tolerance. *Transplantation* 87(9 Suppl.):S55–57; 2009.
- Bartholomew, A.; Sturgeon, C.; Siatskas, M.; Ferrer, K.; McIntosh, K.; Patil, S.; Hardy, W.; Devine, S.; Ucker, D.; Deans, R.; Moseley, A.; Hoffman, R. Mesenchymal stem cells suppress lymphocyte proliferation in vitro and prolong skin graft survival in vivo. *Exp. Hematol.* 30(1):42–48; 2002.
- Campagnoli, C.; Roberts, I. A.; Kumar, S.; Bennett, P. R.; Bellantuono, I.; Fisk, N. M. Identification of mesenchymal stem/progenitor cells in human first-trimester fetal blood, liver, and bone marrow. *Blood* 98(8):2396–2402; 2001.
- Cao, F.; Lin, S.; Xie, X.; Ray, P.; Patel, M.; Zhang, X.; Drukker, M.; Dylla, S. J.; Connolly, A. J.; Chen, X.; Weissman, I. L.; Gambhir, S. S.; Wu, J. C. In vivo visualization of embryonic stem cell survival, proliferation, and migration after cardiac delivery. *Circulation* 113(7):1005–1014; 2006.
- Chan, J. L.; Tang, K. C.; Patel, A. P.; Bonilla, L. M.; Pierobon, N.; Ponzio, N. M.; Rameshwar, P. Antigen-presenting property of mesenchymal stem cells occurs during a narrow window at low levels of interferon-gamma. *Blood* 107(12):4817–4824; 2006.
- Corcione, A.; Benvenuto, F.; Ferretti, E.; Giunti, D.; Cappiello, V.; Cazzanti, F.; Risso, M.; Gualandi, F.; Mancardi, G. L.; Pistoia, V.; Uccelli, A. Human mesenchymal stem cells modulate B-cell functions. *Blood* 107(1):367–372; 2006.
- Dahlke, M. H.; Hoogduijn, M.; Eggenhofer, E.; Popp, F. C.; Renner, P.; Slowik, P.; Rosenauer, A.; Piso, P.; Geissler, E. K.; Lange, C.; Chabannes, D.; Mazzanti, B.; Bigenzahn, S.; Bertolino, P.; Kunter, U.; Introna, M.; Rambaldi, A.; Capelli, C.; Perico, N.; Casiraghi, F.; Noris, M.; Gotti, E.; Seifert, M.; Saccardi, R.; Versapag, H. W.; van Hoek, B.; Bartholomew, A.; Wekerle, T.; Volk, H. D.; Remuzzi, G.; Deans, R.; Lazarus, H.; Schlitt, H. J.; Baan, C. C. Toward MSC in solid organ transplantation: 2008 position paper of the MISOT study group. *Transplantation* 88(5):614–619; 2009.
- Di Nicola, M.; Carlo-Stella, C.; Magni, M.; Milanesi, M.; Longoni, P. D.; Matteucci, P.; Grisanti, S.; Gianni, A. M. Human bone marrow stromal cells suppress T-lymphocyte proliferation induced by cellular or nonspecific mitogenic stimuli. *Blood* 99(10):3838–3843; 2002.
- Gang, E. J.; Bosnakovski, D.; Figueiredo, C. A.; Visser, J. W.; Perlingeiro, R. C. SSEA-4 identifies mesenchymal stem cells from bone marrow. *Blood* 109(4):1743–1751; 2007.
- Glennie, S.; Soeiro, I.; Dyson, P. J.; Lam, E. W.; Dazzi, F. Bone marrow mesenchymal stem cells induce division arrest anergy of activated T cells. *Blood* 105(7):2821–2827; 2005.
- Horwitz, E. M.; Le Blanc, K.; Dominici, M.; Mueller, I.; Slaper-Cortenbach, I.; Marini, F. C.; Deans, R. J.; Krause, D. S.; Keating, A. Clarification of the nomenclature for MSC: The International Society for Cellular Therapy position statement. *Cytotherapy* 7(5):393–395; 2005.
- Inoue, S.; Popp, F. C.; Koehl, G. E.; Piso, P.; Schlitt, H. J.; Geissler, E. K.; Dahlke, M. H. Immunomodulatory effects of mesenchymal stem cells in a rat organ transplant model. *Transplantation* 81(11):1589–1595; 2006.
- Jiang, X. X.; Zhang, Y.; Liu, B.; Zhang, S. X.; Wu, Y.; Yu, X. D.; Mao, N. Human mesenchymal stem cells inhibit differentiation and function of monocyte-derived dendritic cells. *Blood* 105(10):4120–4126; 2005.
- Kim, B. G.; Hwang, D. H.; Lee, S. I.; Kim, E. J.; Kim, S. U. Stem cell-based cell therapy for spinal cord injury. *Cell Transplant.* 16(4):355–364; 2007.
- Kim, D. W.; Chung, Y. J.; Kim, T. G.; Kim, Y. L.; Oh, I. H. Cotransplantation of third-party mesenchymal stromal cells can alleviate single-donor predominance and increase engraftment from double cord transplantation. *Blood* 103(5):1941–1948; 2004.
- Krampera, M.; Cosmi, L.; Angeli, R.; Pasini, A.; Liotta, F.; Andreini, A.; Santarlasci, V.; Mazzinghi, B.; Pizzolo, G.; Vinante, F.; Romagnani, P.; Maggi, E.; Romagnani, S.; Annunziato, F. Role for interferon-gamma in the immunomodulatory activity of human bone marrow mesenchymal stem cells. *Stem Cells* 24(2):386–398; 2006.
- Krampera, M.; Glennie, S.; Dyson, J.; Scott, D.; Laylor, R.; Simpson, E.; Dazzi, F. Bone marrow mesenchymal stem cells inhibit the response of naive and memory antigen-specific T cells to their cognate peptide. *Blood* 101(9):3722–3729; 2003.
- Le Blanc, K.; Rasmusson, I.; Sundberg, B.; Gotherstrom, C.; Hassan, M.; Uzunel, M.; Ringden, O. Treatment of severe acute graft-versus-host disease with third party haploidentical mesenchymal stem cells. *Lancet* 363(9419):1439–1441; 2004.
- Le Blanc, K.; Tammik, L.; Sundberg, B.; Haynesworth,

- S. E.; Ringden, O. Mesenchymal stem cells inhibit and stimulate mixed lymphocyte cultures and mitogenic responses independently of the major histocompatibility complex. *Scand. J. Immunol.* 57(1):11–20; 2003.
21. Loh, Y. H.; Wu, Q.; Chew, J. L.; Vega, V. B.; Zhang, W.; Chen, X.; Bourque, G.; George, J.; Leong, B.; Liu, J.; Wong, K. Y.; Sung, K. W.; Lee, C. W.; Zhao, X. D.; Chiu, K. P.; Lipovich, L.; Kuznetsov, V. A.; Robson, P.; Stanton, L. W.; Wei, C. L.; Ruan, Y.; Lim, B.; Ng, H. H. The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. *Nat. Genet.* 38(4):431–440; 2006.
 22. Meisel, R.; Zibert, A.; Laryea, M.; Gobel, U.; Daubener, W.; Dilloo, D. Human bone marrow stromal cells inhibit allogeneic T-cell responses by indoleamine 2,3-dioxygenase-mediated tryptophan degradation. *Blood* 103(12):4619–4621; 2004.
 23. Nasef, A.; Chapel, A.; Mazurier, C.; Bouchet, S.; Lopez, M.; Mathieu, N.; Sensebe, L.; Zhang, Y.; Gorin, N. C.; Thierry, D.; Fouillard, L. Identification of IL-10 and TGF-beta transcripts involved in the inhibition of T-lymphocyte proliferation during cell contact with human mesenchymal stem cells. *Gene Expr.* 13(4–5):217–226; 2007.
 24. Nasef, A.; Mathieu, N.; Chapel, A.; Frick, J.; Francois, S.; Mazurier, C.; Boutarfa, A.; Bouchet, S.; Gorin, N. C.; Thierry, D.; Fouillard, L. Immunosuppressive effects of mesenchymal stem cells: involvement of HLA-G. *Transplantation* 84(2):231–237; 2007.
 25. Nauta, A. J.; Fibbe, W. E. Immunomodulatory properties of mesenchymal stromal cells. *Blood* 110(10):3499–3506; 2007.
 26. Nauta, A. J.; Westerhuis, G.; Kruisselbrink, A. B.; Lurvink, E. G.; Willemze, R.; Fibbe, W. E. Donor-derived mesenchymal stem cells are immunogenic in an allogeneic host and stimulate donor graft rejection in a nonmyeloablative setting. *Blood* 108(6):2114–2120; 2006.
 27. Pochampally, R. R.; Smith, J. R.; Ylostalo, J.; Prockop, D. J. Serum deprivation of human marrow stromal cells (hMSCs) selects for a subpopulation of early progenitor cells with enhanced expression of OCT-4 and other embryonic genes. *Blood* 103(5):1647–1652; 2004.
 28. Popp, F. C.; Eggenhofer, E.; Renner, P.; Geissler, E. K.; Piso, P.; Schlitt, H. J.; Dahlke, M. H. Mesenchymal stem cells can affect solid organ allograft survival. *Transplantation* 87(9 Suppl.):S57–62; 2009.
 29. Potian, J. A.; Aviv, H.; Ponzio, N. M.; Harrison, J. S.; Rameshwar, P. Veto-like activity of mesenchymal stem cells: Functional discrimination between cellular responses to alloantigens and recall antigens. *J. Immunol.* 171(7):3426–3434; 2003.
 30. Ramos, G. A.; Hare, J. M. Cardiac cell-based therapy: Cell types and mechanisms of actions. *Cell Transplant.* 16(9):951–961; 2007.
 31. Rasmusson, I. Immune modulation by mesenchymal stem cells. *Exp. Cell Res.* 312(12):2169–2179; 2006.
 32. Riekstina, U.; Cakstina, I.; Parfejevs, V.; Hoogduijn, M.; Jankovskis, G.; Muiznieks, I.; Muceniece, R.; Ancans, J. Embryonic stem cell marker expression pattern in human mesenchymal stem cells derived from bone marrow, adipose tissue, heart and dermis. *Stem Cell Rev.* 5(4):378–386; 2009.
 33. Ringden, O.; Uzunel, M.; Rasmusson, I.; Remberger, M.; Sundberg, B.; Lonnie, H.; Marschall, H. U.; Dlugosz, A.; Szakos, A.; Hassan, Z.; Omazic, B.; Aschan, J.; Barkholt, L.; Le Blanc, K. Mesenchymal stem cells for treatment of therapy-resistant graft-versus-host disease. *Transplantation* 81(10):1390–1397; 2006.
 34. Romanov, Y. A.; Svintsitskaya, V. A.; Smirnov, V. N. Searching for alternative sources of postnatal human mesenchymal stem cells: Candidate MSC-like cells from umbilical cord. *Stem Cells* 21(1):105–110; 2003.
 35. Romieu-Mourez, R.; Francois, M.; Boivin, M. N.; Stagg, J.; Galipeau, J. Regulation of MHC class II expression and antigen processing in murine and human mesenchymal stromal cells by IFN-gamma, TGF-beta, and cell density. *J. Immunol.* 179(3):1549–1558; 2007.
 36. Sobiesiak, M.; Sivasubramanian, K.; Hermann, C.; Tan, C.; Orgel, M.; Trembl, S.; Cerabona, F.; Dezwart, P.; Ochs, U.; Muller, C. A.; Gargett, C. E.; Kalbacher, H.; Buhring, H. J. The mesenchymal stem cell antigen MSCA-1 is identical to tissue non-specific alkaline phosphatase. *Stem Cells Dev.* 19(5):669–677; 2009.
 37. Sotiropoulou, P. A.; Perez, S. A.; Gritzapis, A. D.; Baxevanis, C. N.; Papamichail, M. Interactions between human mesenchymal stem cells and natural killer cells. *Stem Cells* 24(1):74–85; 2006.
 38. Stagg, J.; Pommey, S.; Eliopoulos, N.; Galipeau, J. Interferon-gamma-stimulated marrow stromal cells: A new type of nonhematopoietic antigen-presenting cell. *Blood* 107(6):2570–2577; 2006.
 39. Sundin, M.; Barrett, A. J.; Ringden, O.; Uzunel, M.; Lonnie, H.; Dackland, A. L.; Christensson, B.; Blanc, K. L. HSCT recipients have specific tolerance to MSC but not to the MSC donor. *J. Immunother.* 32(7):755–764; 2009.
 40. Torrente, Y.; Polli, E. Mesenchymal stem cell transplantation for neurodegenerative diseases. *Cell Transplant.* 17(10–11):1103–1113; 2008.
 41. Tse, W. T.; Pendleton, J. D.; Beyer, W. M.; Egalka, M. C.; Guinan, E. C. Suppression of allogeneic T-cell proliferation by human marrow stromal cells: Implications in transplantation. *Transplantation* 75(3):389–397; 2003.
 42. Uccelli, A.; Moretta, L.; Pistoia, V. Immunoregulatory function of mesenchymal stem cells. *Eur. J. Immunol.* 36(10):2566–2573; 2006.
 43. Weiss, M. L.; Anderson, C.; Medicetty, S.; Seshareddy, K. B.; Weiss, R. J.; VanderWerff, I.; Troyer, D.; McIntosh, K. R. Immune properties of human umbilical cord Wharton's jelly-derived cells. *Stem Cells* 26(11):2865–2874; 2008.
 44. Wilson, K. D.; Huang, M.; Wu, J. C. Bioluminescence reporter gene imaging of human embryonic stem cell survival, proliferation, and fate. *Methods Mol. Biol.* 574:87–103; 2009.
 45. Yanez, R.; Lamana, M. L.; Garcia-Castro, J.; Colmenero, I.; Ramirez, M.; Bueren, J. A. Adipose tissue-derived mesenchymal stem cells have in vivo immunosuppressive properties applicable for the control of the graft-versus-host disease. *Stem Cells* 24(11):2582–2591; 2006.
 46. Zappia, E.; Casazza, S.; Pedemonte, E.; Benvenuto, F.; Bonanni, I.; Gerdoni, E.; Giunti, D.; Ceravolo, A.; Cazzanti, F.; Frassoni, F.; Mancardi, G.; Uccelli, A. Mesenchymal stem cells ameliorate experimental autoimmune encephalomyelitis inducing T-cell anergy. *Blood* 106(5):1755–1761; 2005.
 47. Zhang, J.; Li, Y.; Chen, J.; Cui, Y.; Lu, M.; Elias, S. B.; Mitchell, J. B.; Hammill, L.; Vanguri, P.; Chopp, M. Human bone marrow stromal cell treatment improves neuro-

- logical functional recovery in EAE mice. *Exp. Neurol.* 195(1):16–26; 2005.
48. Zhang, W.; Ge, W.; Li, C.; You, S.; Liao, L.; Han, Q.; Deng, W.; Zhao, R. C. Effects of mesenchymal stem cells on differentiation, maturation, and function of human monocyte-derived dendritic cells. *Stem Cells Dev.* 13(3): 263–271; 2004.
49. Zhang, Z. Y.; Teoh, S. H.; Chong, M. S.; Schantz, J. T.; Fisk, N. M.; Choolani, M. A.; Chan, J. Superior osteogenic capacity for bone tissue engineering of fetal compared with perinatal and adult mesenchymal stem cells. *Stem Cells* 27(1):126–137; 2009.
50. Zhou, H. P.; Yi, D. H.; Yu, S. Q.; Sun, G. C.; Cui, Q.; Zhu, H. L.; Liu, J. C.; Zhang, J. Z.; Wu, T. J. Administration of donor-derived mesenchymal stem cells can prolong the survival of rat cardiac allograft. *Transplant. Proc.* 38(9):3046–3051; 2006.
51. Zuk, P. A.; Zhu, M.; Ashjian, P.; De Ugarte, D. A.; Huang, J. I.; Mizuno, H.; Alfonso, Z. C.; Fraser, J. K.; Benhaim, P.; Hedrick, M. H. Human adipose tissue is a source of multipotent stem cells. *Mol. Biol. Cell* 13(12): 4279–4295; 2002.

